Recombinant Human OX-2/MOX1/CD200 Protein (Fc Tag) 🤷

Catalog Number: PKSH032841



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	OX-2 Membrane Glycoprotein;CD200;MOX1;MOX2	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Gln31-Gly232	
Accession	P41217	
Calculated Molecular Weight	49.5 kDa	
Observed molecular weight	65-90 kDa	
Tag	C-Fc	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

## Data

kDa	MK	R
120 90		Companya in
60		and the second second
40		
30	-	
20	-	1. m.
14	Land .	

> 95 % as determined by reducing SDS-PAGE.

## Background

CD200 is a transmembrane immunoregulatory protein that belongs to the immunoglobulin superfamily. It contains one Ig like V type domain and one Ig like C2 type domain in its extracelluar domain. CD200 is widely but not ubiquitously expressed. Its receptor (CD200R) is restricted primarily to mast cells; basophils; macrophages; and dendritic cells; which suggests myeloid cell regulation as the major function of CD200. CD200 and CD200R associate via their respective N-terminal Ig-like domains. In myeloid cells; CD200R initiates inhibitory signals following receptor-ligand contact. In T cells; CD200 functions as a co-stimulatory molecule independent of the CD28 pathway. In addition; CD200 also plays an important role in prevention of graft rejection; autoimmune diseases and spontaneous abortion.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com