

Recombinant Human PPCDC Protein (His Tag)

Catalog Number:PKSH032896



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Phosphopantothenoylcysteine Decarboxylase;PPC-DC;PPCDC;COAC
Species	Human
Expression Host	E.coli
Sequence	Met 1-Ser204
Accession	Q96CD2
Calculated Molecular Weight	24.6 kDa
Observed molecular weight	27 kDa
Tag	N-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 50mM NaCl, 1mM DTT, 10% Glycerol, pH 8.0.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Phosphopantothenoylcysteine Decarboxylase (PPC-DC) is an essential enzyme in the biosynthesis of Coenzyme A and catalyzes the decarboxylation of PPC to Phosphopantetheine. PPC-DC catalyzes the decarboxylation of the Cysteine moiety of 4-Phosphopantothenoylcysteine (PPC) to form 4-Phosphopantetheine (PPantSH), this reaction forms part of the biosynthesis of Coenzyme A. The enzyme is a member of the larger family of Cysteine Decarboxylases including the Lantibiotic-Biosynthesizing enzymes EpiD and MrsD, all of which use a tightly bound Flavin cofactor to oxidize the Thiol moiety of the substrate to a Thioaldehyde.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017