

## Recombinant Human Protocadherin-1/PCDH1 Protein (His Tag)

Catalog No. PKSH032977

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                                    |   |
|------------------------------------|---|
| <b>Synonyms</b>                    | Protocadherin-1; Cadherin-Like Protein 1; Protocadherin-42; PC42; PCDH1 |
| <b>Species</b>                     | Human   |
| <b>Expression Host</b>             | HEK293 Cells  |
| <b>Sequence</b>                    | Thr58-Asn852  |
| <b>Accession</b>                   | Q08174  |
| <b>Calculated Molecular Weight</b> | 87.3 kDa  |
| <b>Observed molecular weight</b>   | 120 kDa   |
| <b>Tag</b>                         | C-His   |
| <b>Bioactivity</b>                 | Not validated for activity  |

### Properties

|                       |  |
|-----------------------|--|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.   |
| <b>Endotoxin</b>      | < 1.0 EU per µg of the protein as determined by the LAL method.  |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.                      |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.  |
| <b>Formulation</b>    | Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.   |

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Protocadherin-1, also known as Cadherin-like protein 1, Protocadherin-42 and PCDH1, belongs to the protocadherin subfamily within the cadherin superfamily. PCDH1 contains seven cadherin-like domains, a transmembrane region and a

### For Research Use Only

C-terminal cytoplasmic region. PCDH1 can be detected as early as embryonic day 9.5. In early embryogenesis, expression is especially prominent in blood vessels. The tight spatial and temporal regulation of Pcdh1 expression suggests that this protocadherin plays multiple roles not only during development but also in mature tissues and organs. In addition, protocadherin-1 is involved in cell-cell interaction processes and in neural cell adhesion.