

Recombinant Human SerpinG1/C1IN Protein (His Tag)

Catalog No. PKSH033043

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Plasma Protease C1 Inhibitor;C1 Inh;C1Inh;C1 Esterase Inhibitor;C1-Inhibiting Factor;Serpin G1;SERPING1;C1IN;C1NH;HAE1;HAE2
Species	Human
Expression Host	HEK293 Cells
Sequence	Asn23-Ala500
Accession	AAH11171.1
Calculated Molecular Weight	53.9 kDa
Observed molecular weight	102 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

As protease inhibitors, serpins have an array of functions including regulating blood clotting, the complement pathway, extracellular matrix remodeling, and cell motility. Serpin G1 is a serine protease inhibitor protein. It is the largest member among the serpin class of proteins. Remarkably, Serpin G1 has a 2-domain structure, unlike most family members. The C-terminal serpin domain is similar to other serpins, and this part of Serpin G1 provides the inhibitory activity. The N-terminal domain is not essential for Serpin G1 to inhibit proteinases and has no similarity to other proteins. The main function of Serpin G1 is the inhibition of the complement system to prevent spontaneous activation. Serpin G1 is an acute phase protein and circulates in blood at levels of around 0.25g/L, whose levels rise 2-fold during inflammation. Although named after its complement inhibitory activity, Serpin G1 also inhibits proteinases of the fibrinolytic, clotting, and kinin pathways. Most notably, Serpin G1 play a potentially crucial role in regulating important physiological pathways including complement activation, blood coagulation, fibrinolysis and the generation of kinins. It is also the most important physiological inhibitor of fXIIa, chymotrypsin and plasma kallikrein.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017