

Recombinant Human SIGLEC9/CD329 Protein (Fc Tag)

Catalog No. PKSH033055

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Sialic acid-binding Ig-like lectin 9;Siglec-9;CDw329;Protein FOAP-9;SIGLEC9
Species	Human
Expression Host	HEK293 Cells
Sequence	Gln18-Gly348
Accession	AAH35365.2
Calculated Molecular Weight	63.2 kDa
Observed molecular weight	90-120 kDa
Tag	C-Fc
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Sialic acid-binding Ig-like lectin 9(Siglec 9) is expressed by peripheral blood leukocytes (neutrophils and monocytes but not eosinophils); and found in liver; fetal liver; bone marrow; placenta; spleen and in lower levels in skeletal muscle; fetal

For Research Use Only

brain and so on. It is a putative adhesion molecule that mediates sialic-acid dependent binding to cells. It also binds to alpha-2;3- or alpha-2;6-linked sialic acid. The sialic acid recognition site may be masked by cis interactions with sialic acids on the same cell surface.