

Recombinant Human STAT3 Protein (His Tag)

Catalog No. PKSH033057

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Signal Transducer and Activator of Transcription 3;Acute-Phase Response Factor;STAT3;APRF
Species	Human
Expression Host	E.coli
Sequence	Met 1-Asn175
Accession	P40763
Calculated Molecular Weight	21.8 kDa
Observed molecular weight	19&46 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 10% Trehalose, 50mM NaCl, 0.05% Tween 80, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Signal Transducer and Activator of Transcription 3 (STAT3) belongs to the transcription factor STAT family. STAT3 contains one SH2 domain and is a transcription factor expressed in most cell types. STAT3 is activated by multiple cytokines and growth factors including: IFN- α , IL-10, IL-6, IL-11, IL-12, IL-2, EGF etc. STAT3 functions as signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF and other growth factors. In addition, STAT3 may also mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and FGFR4.