A Reliable Research Partner in Life Science and Medicine

# **Recombinant Human STAT5B Protein (His Tag)**

Catalog No. PKSH033058

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Synonyms Signal Transducer and Activator of Transcription 5B;STAT5B

Species Human
Expression Host E.coli

Sequence Met 1-Thr321

AccessionP51692Calculated Molecular Weight38.4 kDaObserved molecular weight36 kDaTagC-His

**Bioactivity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

**Shipping** This product is provided as liquid. It is shipped at frozen temperature with blue


ice/gel packs. Upon receipt, store it immediately at < - 20°C.

**Formulation** Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 10% Trehalose, 1mM

DTT, 0.05% Tween 80, pH 8.5.

**Reconstitution** Not Applicable

### Data



> 95 % as determined by reducing SDS-PAGE.

### **Background**

Signal Transducer and Activator of Transcription 5b (STAT5B) is a member of the STAT family of transcription factors. They are responsible for an array of cellular activities including regulating growth, survival, differentiation, motility, and the immune response. STAT5B mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different growth hormones. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. Signal transduction and

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

activator of transcription 5 (STAT5) is a member of the Jak/STAT signal transduction pathway and is activated by a variety of cytokines (IL22, IL6). STAT5 has two isoforms (A and B) that share 93% amino acid identity and bind the DNA consensus site TTCN3GAA. STAT5 mediates cytokine signaling by acting as a signal transducer in the cytoplasm and, upon phosphorylation, translocates to the nucleus and activates transcription of specific genes. STAT5 is involved in a wide array of biological processes ranging from regulating apoptosis to adult mammary gland proliferation, differentiation and survival.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com