# Recombinant Human SLAMF5/CD84 Protein (aa 22-220, His Tag)



Catalog Number: PKSH033062

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

Synonyms SLAM family member 5;Cell surface antigen MAX.3;Hly9-beta;Leukocyte

differentiation antigen CD84; Signaling lymphocytic activation molecule

5;CD84;SLAMF5;hCD84;LY9B;mCD84

Species Human

Expression Host HEK293 Cells
Sequence Lys22-Arg220
Accession Q9UIB8
Calculated Molecular Weight 23.1 kDa
Observed molecular weight 35-43 kDa
Tag C-His

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

# **Background**

SLAM family member 5 (SLAMF5/CD84) is a type I transmembrane protein in the SLAM subgroup of the CD2 family. SLAM family proteins regulate multiple aspects of immune system function. Mature human CD84 consists of a 204 amino acid (aa) extracellular domain (ECD) with two Iglike domains, a 21 aa transmembrane segment, and a 99 aa cytoplasmic domain with two immunoreceptor tyrosinebased switch motifs (ITSMs). CD84 exhibits homophilic binding which is mediated by the N-terminal Ig-like domain. Ligation induces tyrosine phosphorylation in the cytoplasmic ITSMs which then recruit the signaling adaptor molecules SAP (SLAM-associated protein) and EAT-2(EWS/Fli1-activated

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

# Recombinant Human SLAMF5/CD84 Protein (aa 22-220, His Tag)



Catalog Number:PKSH033062

transcript 2).CD84 signaling inhibits Fc epsilon RI-induced mast cell activation but enhances platelet activation. LPS-induced macrophage activation,T cell proliferation and IFN- $\gamma$ production, and the interactions between T cells and B cells that are required for germinal center formation.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>