
## Recombinant Human SLAMF6/Ly108 Protein (aa 22-225, Fc Tag)



Catalog Number: PKSH033063

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description                 |  |
|-----------------------------|--|
| Synonyms                    | SLAM Family Member 6;Activating NK Receptor;NK-T-B-Antigen;NTB-<br>A;CD352;SLAMF6;KALI;Ly108;NTBA;SF2000   |
| Species                     | Human  |
| Expression Host             | HEK293 Cells   |
| Sequence                    | Gln22-Lys225   |
| Accession                   | Q96DU3   |
| Calculated Molecular Weight | 49.6 kDa   |
| Observed molecular weight   | 55-65 kDa  |
| Tag                         | C-Fc   |
| Properties                  |  |
| Purity                      | > 90 % as determined by reducing SDS-PAGE.   |
| Endotoxin                   | < 1.0 EU per $\mu$ g of the protein as determined by the LAL method.   |
| Storage                     | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.                |
| Shipping                    | This product is provided as lyophilized powder which is shipped with ice packs.  |
| Formulation                 | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as<br>protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual. |
| Reconstitution              | Please refer to the printed manual for detailed information.   |
| Data                        |  |



> 90 % as determined by reducing SDS-PAGE.

### Background

SLAM Family Member 6 (SLAMF6) is a single-pass type I membrane protein that belongs to the SLAM subgroup of the CD2 family. Human SLAMF6/ NTB-A contains a 205 amino acid extracellular domain (ECD) with one Ig-like V-set and one Ig-like C2-set domain; a 21 amino acid transmembrane segment and an 84 amino acid cytoplasmic domain; with two immunoreceptor tyrosine-based switch motifs. SLAMF6 is a homodimer. SLAMF6 can interact with PTN6 and; upon phosphorylation; with PTN11 and SH2D1A/SAP. Phosphorylation-dependent NTB-A association with SAP is required for full production of IFN-γ by NK cells and independent of EAT-2 binding. It Triggers cytolytic activity only in natural killer cells (NK) expressing high surface densities of natural cytotoxicity receptors. On B cells; NTB-A modulates

### For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

# Recombinant Human SLAMF6/Ly108 Protein (aa 22-225, Fc Tag)



Catalog Number:PKSH033063

immunoglobulin class switching and the balance between tolerance and autoimmunity.

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com