

## Recombinant Human TFPI2 Protein (His Tag)

**Catalog No.** PKSH033117

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	Tissue Factor Pathway Inhibitor 2;TFPI-2;Placental Protein 5;PP5;TFPI2
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Asp23-Lys213
<b>Accession</b>	P48307
<b>Calculated Molecular Weight</b>	22.9 kDa
<b>Observed molecular weight</b>	18-33 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Human Tissue Factor Pathway Inhibitor 2 (TFPI2) has a N-terminal acidic region, three Kunitz domains separated with by two linker regions, and a C-terminal basic region. TFPI2 has the function of regulating plasmin-mediated matrix

### For Research Use Only

remodeling, inhibits trypsin, plasmin, factor VIIa/tissue factor and weakly factor Xa.. TFPI2 has no effect on thrombin. TFPI2 may contribute to tumor progression in many cancers; in these cancers, the expression of TFPI2 can be down-regulated.