

Recombinant Human TMX2/TXNDC14 Protein (His Tag)

Catalog No. PKSH033123

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Thioredoxin-related transmembrane protein 2;Cell proliferation-inducing gene 26 protein;Thioredoxin domain-containing protein 14;TMX2
Species	Human
Expression Host	E.coli
Sequence	Met 125-Lys296
Accession	Q9Y320
Calculated Molecular Weight	21.9 kDa
Observed molecular weight	23 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

TMX2 is a single-pass type I membrane protein and contains 1 thioredoxin domain. Thioredoxin plays an important role in various cellular processes through redox regulation. The molecular cloning and characterization of one member of the thioredoxin superfamily, designated as TMX2. The TMX2 cDNA consists of 1644 nucleotides and contains an open reading frame encoding a protein of 372 amino acids with a predicted molecular mass of 42.5 kDa and an isoelectric point of 8.94. The TMX2 protein may possess an N-terminal signal peptide, a potential transmembrane domain, an Myb DNA-binding domain repeat signature, a thioredoxin consensus pattern, an endoplasmic reticulum (ER) membrane retention signal (KKXX-like motif), and a dileucine motif in the tail.