

# Recombinant Human YY1 Protein (His Tag)

Catalog Number:PKSH033137



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Transcriptional repressor protein YY1;Delta transcription factor;INO80 complex subunit S;NF-E1;Yin and yang 1;INO80S
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Val221-Gly321
<b>Accession</b>	P25490
<b>Calculated Molecular Weight</b>	12.6 kDa
<b>Observed molecular weight</b>	19 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Transcriptional repressor protein YY1 contains 4 C2H2-type zinc fingers and belongs to the YY transcription factor family. Multifunctional transcription factor exhibits positive and negative control on a large number of cellular and viral genes by binding to sites overlapping the transcription start site. The effect on transcription regulation of the protein is depending upon the context in which it binds and diverse mechanisms of action include direct activation or repression, indirect activation or repression via cofactor recruitment, or activation or repression by disruption of binding sites or conformational DNA changes. Its activity is regulated by transcription factors and cytoplasmic proteins that have been shown to abrogate or completely inhibit YY1-mediated activation or repression.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017