

Recombinant Human TGFB3 Protein

Catalog No. PKSH033140

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Transforming growth factor beta-3;TGFB3;TGF-beta-3;Latency-associated peptide;LAP
Species	Human
Expression Host	HEK293 Cells
Sequence	Ala301-Ser412(Tyr340Phe)
Accession	P10600
Calculated Molecular Weight	12.7 kDa
Observed molecular weight	12-14 kDa
Tag	None
Bioactivity	Measured by its ability to inhibit the IL-4-dependent proliferation of TF-1 mouse T cells. The ED50 for this effect is 10-80 pg/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.01 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 50mM Glycine-HCl, 150mM NaCl, pH 2.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

Transforming growth factor beta 3 (TGFB3) is a member of a TGF- β superfamily which is defined by their structural and functional similarities. TGFB3 is secreted as a complex with LAP. This latent form of TGFB3 becomes active upon cleavage by plasmin, matrix metalloproteases, thrombospondin -1, and a subset of integrins. It binds with high affinity to TGF- β RII, a type II serine/threonine kinase receptor. TGFB3 is involved in cell differentiation, embryogenesis and development. It is believed to regulate molecules involved in cellular adhesion and extracellular matrix (ECM) formation during the process of palate development. Without TGF- β 3, mammals develop a deformity known as a cleft palate.

For Research Use Only