A Reliable Research Partner in Life Science and Medicine

Recombinant Human Tie1 Protein (His Tag)

Catalog No. PKSH033171

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Tyrosine-Protein Kinase Receptor Tie-1;TIE1;TIE

Species Human

Expression HostHEK293 CellsSequenceAla22-Gln760

AccessionP35590Calculated Molecular Weight81.0 kDaObserved molecular weight87 kDaTagC-His

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 500mM NaCl, 10%

Sucrose, pH8.2.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

<u>Data</u>



> 95 % as determined by reducing SDS-PAGE.

Background

TIE-1 (Tyrosine Kinase with Ig and EGF Homology domains 1) and TIE-2/Tek comprise a receptor tyrosine kinase

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

(RTK) subfamily. These receptors are expressed on endothelial and hematopoietic progenitor cells and play critical roles in angiogenesis, vasculogenesis and hematopoiesis. Human TIE-1 cDNA encodes a 1124 amino acid (aa) residue precursor protein with an 18aa signal peptide, a 727 aa extracellular domain and a 354 aa cytoplasmic domain. so far, two ligands have been described for TIE-2 [angiopoietin-1 (Ang1) and angiopoietin-2 (Ang2)], but no ligand was found for TIE-1.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com