

Recombinant Human UROS/UROIIIIS Protein (His Tag)

Catalog No. PKSH033200

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Uroporphyrinogen-III Synthase;UROIIIIS;UROS;Hydroxymethylbilane Hydrolyase [Cyclizing];Uroporphyrinogen-III Cosynthase;UROS
Species	Human
Expression Host	E.coli
Sequence	Met 1-Cys265
Accession	P10746
Calculated Molecular Weight	29.7 kDa
Observed molecular weight	29 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 100mM NaCl, 10% Glycerol, pH 8.0.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Uroporphyrinogen-III Synthase is an enzyme which belongs to the uroporphyrinogen-III synthase family. Uroporphyrinogen-III Synthase is ubiquitous and it is involved in Porphyrin metabolism. Porphyrins act as cofactors for a multitude of enzymes that perform a variety of processes within the cell such as Methionine synthesis (Vitamin B12) or oxygen transport (Heme). Uroporphyrinogen-III Synthase can catalyze cyclization of the linear Tetrapyrrole,

For Research Use Only

Hydroxymethylbilane, to the Macrocyclic Uroporphyrinogen III, the branch point for the various sub-pathways leading to the wide diversity of Porphyrins. Defects in Uroporphyrinogen-III Synthase are the cause of Congenital Erythropoietic Porphyria (CEP).