A Reliable Research Partner in Life Science and Medicine

Recombinant Human Vitronectin/VTN Protein (His Tag)

Catalog No. PKSH033218

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Vitronectin; VN; S-Protein; Serum-Spreading Factor; V75; VTN

Species Human

Expression Host
Sequence
Asp20-Leu478
Accession
AAH05046.1
Calculated Molecular Weight
Observed molecular weight
Tag
HEK293 Cells
Asp20-Leu478
AAH05046.1
Calculated Molecular Weight
C3.4 kDa
60-80 kDa
C-His

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH

8.0.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

<u>Data</u>



> 95 % as determined by reducing SDS-PAGE.

Background

Human Vitronectin/VTN is a cell adhesion and spreading factor. It can be found in the blood and the extracellular matrix

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

(ECM). Vitronectin interacts with glycosaminoglycans and proteoglycans. The multimeric Vitronectin can efficiently bind to and incorporate into the ECM; Vitronectin can support cell adhesion through binding to various integrins and other proteoglycans. Vitronectin can be recognized by certain members of the integrin family and serves as a cell-to-substrate adhesion molecular. It can as a inhibitor of the membrane-damaging effect of the terminal cytolytic complement pathway. Vitronectin contains an endogenous cleavage site; plus cleavage sites for elastase; thrombin; and plasmin.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com