

Recombinant Human VSIG4 Protein (Fc Tag)

Catalog No. PKSH033219

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	V-set and immunoglobulin domain-containing protein 4;VSIG4;Protein Z39Ig;Z39IG;CRIg
Species	Human
Expression Host	HEK293 Cells
Sequence	Arg20-Val284
Accession	Q9Y279
Calculated Molecular Weight	56.3 kDa
Observed molecular weight	75 kDa
Tag	C-Fc
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

V-set and immunoglobulin domain-containing protein 4(VSIG4) is a transmembrane protein contains a signal peptide, a V-

For Research Use Only

type Ig-like domain, a C2-type Ig-like domain, several potential O-glycosylation sites, and an intracellular domain with 2 potential phosphorylation sites and is structurally related to the B7 family of immune regulatory proteins. This protein is also a receptor for the complement component 3 fragments C3b and iC3b. The main function is strong negative regulator of T-cell proliferation and IL2 production and it is also potent inhibitor of the alternative complement pathway convertases. It abundantly expressed in several fetal tissues such as adult tissues, highest expression in lung and placenta and it also expressed in resting macrophages.