

Recombinant Rat CLPS/Colipase Protein (His Tag)

Catalog No. PKSR030146

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CLPS;Colipase
Species	Rat
Expression Host	HEK293 Cells
Sequence	Met1-Gln112
Accession	NP_037271.1
Calculated Molecular Weight	11.9 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Colipase belongs to the colipase family. Structural studies of the complex and of colipase alone have revealed the functionality of its architecture. It is a small protein with five conserved disulphide bonds. Structural analogies have been recognised between a developmental protein, the pancreatic lipase C-terminal domain, the N-terminal domains of

For Research Use Only

lipoxygenases and the C-terminal domain of alpha-toxin. Colipase can only be detected in pancreatic acinar cells, suggesting regulation of expression by tissue-specific elements. Colipase allows lipase to anchor noncovalently to the surface of lipid micelles, counteracting the destabilizing influence of intestinal bile salts. Without colipase the enzyme is washed off by bile salts, which have an inhibitory effect on the lipase. Colipase is a cofactor needed by pancreatic lipase for efficient dietary lipid hydrolysis. It binds to the C-terminal, non-catalytic domain of lipase, thereby stabilising as active conformation and considerably increasing the overall hydrophobic binding site.