

# Recombinant Rat Layilin/LAYN Protein

Catalog Number:PKSR030170



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	LAYN
<b>Species</b>	Rat
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met1-Glu224
<b>Accession</b>	NP_001178926.1
<b>Calculated Molecular Weight</b>	23.7 kDa
<b>Observed molecular weight</b>	34 kDa
<b>Tag</b>	None

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Layilin, a recently characterized as a 55 kDa transmembrane protein with homology to C-type lectins, is present in numerous cell lines and tissue extracts. As one of the adaptor proteins, talin mediates the interactions between the actin filaments and the cell membrane by binding to integral membrane proteins and to the cytoskeleton. Layilin is a newly identified membrane-binding site for talin in peripheral ruffles of spreading cells, a ten-amino acid motif in the layilin cytoplasmic domain is sufficient for talin binding, and its adjacent LH2-LH3 tandem arrays in the cytoplasmic domain provide docking sites for talin. Layilin binds specifically to hyaluronan (HA) through its extracellular domain, a ubiquitous extracellular matrix component in most animal tissues and body fluids, but not to other tested glycosaminoglycans. The research suggests that layilin may mediate signals from extracellular matrix to the cell

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Rat Layilin/LAYN Protein

Catalog Number:PKSR030170



cytoskeleton via interaction with different intracellular binding partners and thereby be involved in the modulation of cortical structures in the cell. All the above actions reveal an interesting parallel between layilin and the known HA receptor CD44. In addition, merlin and radixin have been identified as different intracellular binding partners of layilin. Accordingly, it has been suggested that layilin plays roles in a variety of cellular processes, including cell shape, adhesion, motility, and homeostasis, as well as signal transduction. In addition, layilin might play an important role in the process of invasion and lymphatic metastasis of lung carcinoma.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017