A Reliable Research Partner in Life Science and Medicine

Recombinant Rat CLM-9/TREM4 Protein (His Tag)

Catalog No. PKSR030178

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms CD300LG

Species Rat

Expression Host HEK293 Cells
Sequence Met1-Arg160
Accession XP_002724611.1

Calculated Molecular Weight 17.5 kDa
Observed molecular weight 28 kDa
Tag C-His

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CLM-9, also known as TREM4, is a receptor which belongs to the TREM family. The TREM family of receptors regulates the activity of various cell types of the immune system including neutrophils, monocyte/macrophages, microglia,

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>





A Reliable Research Partner in Life Science and Medicine

and dendritic cells. CLM-9 may mediate L-selectin-dependent lymphocyte rollings. It binds SELL in a calcium dependent manner. CLM-9 also binds lymphocyte which suggests that it functions in lymphocyte adhesion. The major CLM-9 transcript is expressed highly in human heart, skeletal muscle, and placenta. The mouse protein has been shown to be expressed in capillary endothelial cells. Human CLM-9 mediates the uptake of human IgA2 and mouse IgM.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com