

Recombinant Rat CLEC4A3 Protein (Fc Tag)

Catalog No. PKSR030188

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|----------------------------|
| Synonyms | CLEC4A3 |
| Species | Rat |
| Expression Host | HEK293 Cells |
| Sequence | Leu68-Leu237 |
| Accession | Q5YIS0 |
| Calculated Molecular Weight | 48.3 kDa |
| Observed molecular weight | 58 kDa |
| Tag | N-hFc |
| Bioactivity | Not validated for activity |

Properties

| | |
|-----------------------|---|
| Purity | > 85 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 85 % as determined by reducing SDS-PAGE.

Background

CLEC4A3 contains 1 C-type lectin domain and belongs to the C-type lectin-like domain-containing (CLEC) family. Lectins are proteins that are able to recognize and bind with specific carbohydrate molecules. C-type lectins are an

For Research Use Only

important group of proteins found in the immune system of animals. These lectins are named C-type because of their calcium dependent carbohydrate recognition domain (CRD). In the immune system, C-type lectins act as recognition molecules by binding to foreign microorganisms. They also promote the movement and selective adhesion of white blood cells. The C-type lectin has a three-dimensional fold, the CRD, in which calcium ions contribute to the lectin's ability to recognize and bind carbohydrates. In the immune system, carbohydrate recognition contributes to the ability of immune cells to move from one area of the body to another. It also allows immune cells to identify and discriminate between proteins that belong to the host and those that belong to foreign organisms. There are a number of different C-type lectin subfamilies, including collectins, selectins, proteoglycans, and lymphocyte lectins.