
Recombinant Rat LAG3/CD223 Protein (His Tag)

Catalog No. PKSR030201

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	LAG3
Species	Rat
Expression Host	HEK293 Cells
Sequence	Met1-Leu442
Accession	Q5BK54
Calculated Molecular Weight	47.1 kDa
Observed molecular weight	55 kDa
Tag	C-His
Bioactivity	Not validated for activity
Properties	
Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20 mM NaAc, 500 mM NaCl, 10 % Glycerol, 0.02 % tween 20, pH 5.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



> 90 % as determined by reducing SDS-PAGE.

Background

LAG3, also known as CD223 and Lymphocyte activation gene 3, belongs to immunoglobulin (Ig) superfamily. LAG3

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

Elabscience®

contains 4 extracellular Ig-like domains. The LAG3 gene contains 8 exons. It is selectively expressed in activated T and NK cells. LAG3 has a negative regulatory function in T cells. It also acts as as a new marker of T cell induced B cell activation. As a soluble molecule, LAG3 activates antigen-presenting cells through MHC class II signalling, leading to increased antigen-specific T-cell responses in vivo.

For Research Use Only