
## Recombinant Rat CD79B/B29 Protein (His Tag)

### Catalog No. PKSR030227

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description                 |   |
|-----------------------------|---|
| Synonyms                    | CD79B;B29/Ig-beta;Igb   |
| Species                     | Rat   |
| Expression Host             | HEK293 Cells  |
| Sequence                    | Met1-Asp158   |
| Accession                   | O70153  |
| Calculated Molecular Weight | 16.8 kDa  |
| Observed molecular weight   | 28 kDa  |
| Tag                         | C-His   |
| Bioactivity                 | Not validated for activity  |
| Properties                  |   |
| Purity                      | > 90 % as determined by reducing SDS-PAGE.  |
| Endotoxin                   | < 1.0 EU per $\mu$ g of the protein as determined by the LAL method.  |
| Storage                     | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping                    | This product is provided as lyophilized powder which is shipped with ice packs.   |
| Formulation                 | Lyophilized from sterile PBS, pH 7.4<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as<br>protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual.         |
| Reconstitution              | Please refer to the printed manual for detailed information.  |

Data



> 90 % as determined by reducing SDS-PAGE.

## Background

CD79B is a single-pass type I membrane protein. CD79B contains one Ig-like V-type domain and one ITAM domain. CD79B is required in cooperation with CD79A for initiation of the signal transduction cascade activated by the B-cell

#### **For Research Use Only**

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

# **Elabscience**®

antigen receptor complex (BCR), which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79B enhances phosphorylation of CD79A, possibly by recruiting kinases that phosphorylate CD79A or by recruiting proteins that bind to CD79A and protect it from dephosphorylation.

For Research Use Only