Recombinant Rat Semaphorin 4D/SEMA4D/CD100 Protein (Fc Tag)

Catalog No. PKSR030237

Note: Centrifuge before opening to ensure complete recovery of vial contents.

ynonyms	SEMA4D
pecies	Rat
xpression Host	HEK293 Cells
equence	Met1-Met712
ccession	NP_001164034.1
alculated Molecular Weight	103.4 kDa
bserved molecular weight	118 kDa
ag	C-hFc
ioactivity	Not validated for activity
roperties	
urity	> 95 % as determined by reducing SDS-PAGE.
ndotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
C	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
hipping	This product is provided as lyophilized powder which is shipped with ice packs.
	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
econstitution	Please refer to the printed manual for detailed information.

Data

KDa	М
116	
66.2	
45.0	-
35.0	-
25.0	-
18.4 14.4	=

> 95 % as determined by reducing SDS-PAGE.

Background

Semaphorin-4D is also known as A8,BB18, GR3, CD100. Semaphorin-4D belongs to the semaphorin family containing 1 Ig-like C2-type domain, 1 PSI domain and 1 Sema domain. It is the cell surface receptor for PLXN1B and PLXNB2 that

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

plays an important role in cell-cell signaling. It promotes the migration of cerebellar granule cells and of endothelial cells, regulates dendrite and axon branching and morphogenesis. Semaphorin-4D Plays a role in the immune system; Promotes signaling via SRC and PTK2B/PYK2, which then mediates activation of phosphatidylinositol 3-kinase and of the AKT1 signaling cascade.

For Research Use Only