

Recombinant Rat CD63/Tspan-30 Protein (Fc Tag)

Catalog No. PKSR030251

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD63
Species	Rat
Expression Host	HEK293 Cells
Sequence	Ala103-Val203
Accession	P28648
Calculated Molecular Weight	38.1 kDa
Observed molecular weight	42-47 kDa
Tag	N-mFc
Bioactivity	Not validated for activity

Properties

Purity	> 85 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 85 % as determined by reducing SDS-PAGE.

Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune

For Research Use Only

function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 63 (CD63) is a member of the CD family and the transmembrane 4 superfamily, also known as the tetraspanin family. CD63 is a cellular surface glycoprotein characterized by the presence of four hydrophobic domains. CD63 had functions in mediating signal transduction processes and then regulate variety of cellular processes such as cell proliferation, activation and motility. It has reported that CD63 protein associated with tumor progression and served as a blood platelet activation marker and the deficiency of this protein may be associated with Hermansky-Pudlak syndrome.