

Recombinant Rat IL6ST/CD130 Protein (His Tag)

Catalog No. PKSR030390

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Ac1055;Gp130;IL-6rb
Species	Rat
Expression Host	HEK293 Cells
Sequence	Met 4-Glu 618
Accession	NP_001008725.2
Calculated Molecular Weight	68.2 kDa
Observed molecular weight	90-100 kDa
Tag	C-His
Bioactivity	Measured by its ability to inhibit the IL6 R α enhancement of IL6 activity on M1 mouse myeloid leukemia cells. The ED50 for this effect is typically 0.5-2 μ g/ml in the presence of 50 ng/ml recombinant human IL6sR and 100 ng/ml recombinant human IL6.

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

Glycoprotein 130 (also known as gp130, IL6ST, IL6-beta or CD130) is a transmembrane protein which is the founding member of the class of all cytokine receptors. CD130/gp130 is a signal transducer shared by many cytokines, including interleukin 6 (IL6), ciliary neurotrophic factor (CNTF), leukemia inhibitory factor (LIF), and Oncostatin M (OSM). CD130/gp130 functions as a part of the cytokine receptor complex. The activation of this protein is dependent upon the binding of cytokines to their receptors. CD130/gp130 plays a critical role in regulating myocyte apoptosis. Alternatively spliced transcript variants encoding distinct isoforms have been described. A related pseudogene has been identified on chromosome 17. The receptor systems for IL6, LIF, OSM, CNTF, IL11, CTF1 and BSF3 can utilize gp130 for initiating signal transmission. CD130/gp130 binds to IL6/IL6R (alpha chain) complex, resulting in the formation of high-affinity IL6 binding sites, and transduces the signal. CD130/gp130 may have a role in embryonic development. The type I OSM receptor is capable of transducing OSM-specific signaling events.

For Research Use Only