

## Recombinant Mouse CD45/PTPRC Protein

Catalog No. PKSM040296

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	B220;Cd45;CD45R;L-CA;loc;Ly-5;Lyt-4;T200
<b>Species</b>	Mouse
<b>Expression Host</b>	Baculovirus-Insect Cells
<b>Sequence</b>	Arg453-Ser1152
<b>Accession</b>	AAA39458.1
<b>Calculated Molecular Weight</b>	81.0 kDa
<b>Observed molecular weight</b>	93 kDa
<b>Tag</b>	None
<b>Bioactivity</b>	<ol style="list-style-type: none"> <li>1. The specific activity was determined to be 9168 nmol/min/mg using p-nitrophenyl phosphate as substrate.</li> <li>2. Immobilized mouse PTPRC (453-1152) at 10 µg/ml (100 µl/well) can bind biotinylated human Galectin-1 with a linear range of 0.31-2. 5 µg/ml.</li> </ol>

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
<b>Formulation</b>	Supplied as sterile solution of 20mM Tris, 500mM NaCl, 10% glycerol, 3mM DTT, pH 7.4
<b>Reconstitution</b>	Not Applicable

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune

### For Research Use Only

function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Protein tyrosine phosphatase, receptor type C (CD45), also known as PTPRC is a member of the protein tyrosine phosphatase (PTP) family which is known for its function to serve as signaling molecules and to regulate a variety of cellular processes such as cell proliferation, differentiation, mitotic cycle and oncogenic transformation. CD45 is found expression specifically in hemotopietic cells. CD45 consists of an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains. It serves as an essential regulator of T-cell and B-cell antigen receptor signaling through either direct interaction with components of the antigen receptor complex or by activating various Src family kinases required for the antigen receptor signaling and it also can suppress JAK kinases.