

## Recombinant Mouse PTK6/Brk Protein (His & GST Tag)

**Catalog No.** PKSM040298

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	BRK;Sik;tk;Tksk
<b>Species</b>	Mouse
<b>Expression Host</b>	Baculovirus-Insect Cells
<b>Sequence</b>	Met1-Val451
<b>Accession</b>	Q64434
<b>Calculated Molecular Weight</b>	79.8 kDa
<b>Observed molecular weight</b>	66 kDa
<b>Tag</b>	N-His-GST
<b>Bioactivity</b>	The specific activity was determined to be 5 nmol/min/mg using poly [Glu, Tyr] 4:1 as substrate.

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
<b>Formulation</b>	Supplied as sterile solution of 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol
<b>Reconstitution</b>	Not Applicable

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

Tyrosine kinase (PTKs) is a protein that carry out tyrosine phosphorylation, which play a fundamental role in cell proliferation, survival, adhesion, and motility and have also been demenstrated to mediate malignant cell transformation. Overexpression of this protein in mammary epithelial cells leads to sensitization of the cells to epidermal growth factor and results in a partially transformed phenotype. Two classes of PTKs are present in cells: the transmembrane receptor PTKs and the non-receptor PTKs. Tyrosine kinase(PTKs)-6/ BRK is a cytoplasmic non-receptor protein kinase which

### For Research Use Only

may function as an intracellular signal transducer in epithelial tissues. Tyrosine kinase(PTKs)-6/ BRK has been shown to undergo autophosphorylation. It has been found that the constitutive expression of the tyrosine kinase(PTKs)-6/ BRK is in a large proportion of cutaneous T-cell lymphomas and other transformed T- and B-cell populations. State BRK expression was also induced in normal T-cells. In clinical, the cytoplasmic tyrosine kinase PTK6 (BRK) shows elevated expression in approximately two-thirds of primary breast tumours, and is implicated in EGF receptor-dependent signalling and epithelial tumorigenesis.