A Reliable Research Partner in Life Science and Medicine

# Recombinant Mouse SCG3/Secretogranin 3 Protein (His Tag)

Catalog No. PKSM040310

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Synonyms 1B1075;AI385542;Chgd;SgIII

**Species** Mouse

Expression Host HEK293 Cells
Sequence Met1-Leu471
Accession NP\_033156.1
Calculated Molecular Weight 52.4 kDa
Tag C-His

**Bioactivity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

## **Background**

SCG3, also known as secretogranin 3, is a member of the chromogranin/secretogranin family. Members of this family may serve as precursors for biologically active peptides. SCG3 is transported to secretory granules (SGs) in neuroendocrine cells. SCG3 binds strongly to chromogranin A (CgA) in an intragranular milieu and targets CgA to SGs in

### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

 $Email: \underline{tech support@elabscience.com}$ 

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

pituitary and pancreatic endocrine cells. With a sucrose density gradient of rat insulinoma-derived INS-1 cell homogenates, SgIII is localized to the SG fraction and is fractionated to the SG membrane (SGM) despite lacking the transmembrane region.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com