Recombinant Mouse Osteoprotegerin/TNFRSF11B Protein (Fc Tag)

Catalog No. PKSM040324

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Tumor necrosis factor receptor superfamily member 11B;Osteoclastogenesis inhibitory factor;Osteoprotegerin;Tnfrsf11b;Ocif;Opg;TR1
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met1-Leu401
Accession	NP_032790.3
Calculated Molecular Weight	70.2 kDa
Tag	C-hFc
Bioactivity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Osteoprotegerin or TNFRSF11B is a member of the TNF-receptor superfamily. This protein is an osteoblast-secreted decoy receptor that functions as a negative regulator of bone resorption. This protein specifically binds to its ligand,

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

osteoprotegerin ligand, both of which are key extracellular regulators of osteoclast development. Studies of the mouse counterpart also suggest that this protein and its ligand play a role in lymph-node organogenesis and vascular calcification. Alternatively spliced transcript variants of this gene have been reported, but their full length nature has not been determined. Osteoprotegerin/TNFRSF11B acts as decoy receptor for RANKL and thereby neutralizes its function in osteoclastogenesis. This protein may inhibit the activation of osteoclasts and promotes osteoclast apoptosis in vitro. Bone homeostasis seems to depend on the local RANKL/OPG ratio. Osteoprotegerin/TNFRSF11B also play a role in preventing arterial calcification, act as decoy receptor for TRAIL and protect against apoptosis. TRAIL binding blocks the inhibition of osteoclastogenesis.

For Research Use Only