

Recombinant Mouse TROP2/TACSTD2 Protein (Fc Tag)

Catalog Number:PKSM040436



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Tumor-associated calcium signal transducer 2; Tacstd2; Trop2; Cell surface glycoprotein Trop-2; TROP-2; C80403; EGP-1; GA733-1; Ly97
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Gln 270
Accession	Q8BGV3
Calculated Molecular Weight	55.0 kDa
Observed molecular weight	66 kDa
Tag	C-hFc
Bioactivity	Measured by the ability of the immobilized protein to support the adhesion of U937 human histiocytic lymphoma cells. When 5×10^4 cells/well are added to coated plates (40 $\mu\text{g/mL}$, 100 $\mu\text{L/well}$), > 50% cells will adhere after 1 hour at 37°C.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

TROP-2, also referred to as tumor associated calcium signal transducer 2 (TACSTD2), GA733-1 or M1S1, is a cell surface glycoprotein highly expressed in a wide variety of epithelial cancers. In contrast, there is little or no expression of Trop-2 in adult somatic tissue. Because it is a cell surface protein that is selectively expressed in tumor cells, Trop-2 is a potential therapeutic target. The cytoplasmic tail of Trop-2 possesses potential serine and tyrosine phosphorylation sites

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse TROP2/TACSTD2 Protein (Fc Tag)

Catalog Number:PKSM040436



and a phosphatidyl-inositol binding consensus sequence. Trop-2 transduces an intracellular calcium signal, are consistent with the hypothesis that it acts as a cell surface receptor and support a search for a physiological ligand. TROP2 encoding by an intronless gene was originally defined by the monoclonal antibody GA733, and is a member of a family of at least two type I membrane proteins. The other known member is GA733-2, also called EpCAM and TROP1. It has been suggested by studies that the GA733-1 gene was formed by the retroposition of the GA733-2 gene via an mRNA intermediate.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com