

Recombinant Mouse Kallikrein 11/KLK11 Protein (His Tag)

Catalog No. PKSM040438

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	2310015I08Rik;Prss20;TLSP
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Asn 276
Accession	Q9QYN3
Calculated Molecular Weight	27.3 kDa
Observed molecular weight	38-42 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

kallikrein-related peptidase 11 (KLK11), also known as hippostasin, trypsin-like serine protease and PRSS20, is a member of human tissue kallikrein family. It is a subgroup of serine proteases with diverse physiological functions, which

For Research Use Only

is implicated in carcinogenesis and some with potential that serving as novel biomarkers for ovarian and prostate cancer and other diseases. The KLK11 gene is one of the fifteen kallikrein subfamily members located in a cluster on chromosome 19. Two alternatively spliced forms exist, resulting in 250 (isoform 1) and 282 (isoform 2) amino acid sequences. Isoform 2 is identical to isoform 1, except for an inserted 32 amino acid segment. Isoform 1 is predominantly expressed in brain whereas isoform 2 is preferentially expressed in prostate.