Recombinant Mouse Carbonic Anhydrase X/CA10 Protein (His Tag)

Catalog No. PKSM040494

Note: Centrifuge before opening to ensure complete recovery of vial contents.

SpeciesMouseExpression HostHEK293 CellsSequenceMet 1-Asn 300AccessionP61215Calculated Molecular Weight33.1 kDaObserved molecular weight38 kDaTagC-HisBioactivityMeasured by its esterase activity. The specific activity is > 10 pmoles/min/µg.PropertiesPurity> 97 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per µg of the protein as determined by the LAL method.	Description		
Expression HostHEK293 CellsSequenceMet 1-Asn 300AccessionP61215Calculated Molecular Weight33.1 kDaObserved molecular weight38 kDaTagC-HisBioactivityMeasured by its esterase activity. The specific activity is > 10 pmoles/min/µg.PropertiesPurity> 97 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per µg of the protein as determined by the LAL method.	Synonyms	2700029L05Rik;BB085816;Ca10;Car10;RP23-448I9.1	
SequenceMet 1-Asn 300AccessionP61215Calculated Molecular Weight33.1 kDaObserved molecular weight38 kDaTagC-HisBioactivityMeasured by its esterase activity. The specific activity is > 10 pmoles/min/μg.PropertiesPurity> 97 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.	Species	Mouse	
AccessionP61215Calculated Molecular Weight33.1 kDaObserved molecular weight38 kDaTagC-HisBioactivityMeasured by its esterase activity. The specific activity is > 10 pmoles/min/μg.PropertiesPurity> 97 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per µg of the protein as determined by the LAL method.	Expression Host	HEK293 Cells	
Calculated Molecular Weight33.1 kDaObserved molecular weight38 kDaTagC-HisBioactivityMeasured by its esterase activity. The specific activity is > 10 pmoles/min/μg.PropertiesPurity> 97 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per µg of the protein as determined by the LAL method.	Sequence	Met 1-Asn 300	
Observed molecular weight38 kDaTagC-HisBioactivityMeasured by its esterase activity. The specific activity is > 10 pmoles/min/μg.PropertiesPurity> 97 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per µg of the protein as determined by the LAL method.	Accession	P61215	
TagC-HisBioactivityMeasured by its esterase activity. The specific activity is > 10 pmoles/min/μg.PropertiesPurity> 97 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.	Calculated Molecular Weight	33.1 kDa	
BioactivityMeasured by its esterase activity. The specific activity is > 10 pmoles/min/μg.PropertiesPurity> 97 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.	Observed molecular weight	38 kDa	
Properties Purity > 97 % as determined by reducing SDS-PAGE. Endotoxin < 1.0 EU per μg of the protein as determined by the LAL method. Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to	Tag	C-His	
Purity> 97 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.	Bioactivity	Measured by its esterase activity. The specific activity is > 10 pmoles/min/ μ g.	
Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.	Properties		
Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to	Purity	> 97 % as determined by reducing SDS-PAGE.	
	Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
FormulationLyophilized from sterile 25mM Tris, 150mM NaCl, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	Formulation	Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.	
Reconstitution Please refer to the printed manual for detailed information.	Reconstitution	Please refer to the printed manual for detailed information.	

Data

KDa	MK	R
116	-	
66.2	-	
45.0	-	
35.0	-	
25.0	-	
18.4	-	
14.4	-	

> 97 % as determined by reducing SDS-PAGE.

Background

Carbonic anhydrase X, also called carbonic anhydrase - related protein X (CARPX) and CA10, belongs to the CA family of zinc metalloenzymes which catalyze the reversible hydration of carbon dioxide in various biological processes such as

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

respiration, renal tubular acidification and bone resorption. The secreted protein CARPX without CA activity (hydration of CO2) is identified as an acatalytic member of the alpha-carbonic anhydrase subgroup. CARP X expression is detected in the adult total brain and almost all parts of the central nervous system, but not in the fetal brain. Accordingly, CARP X is suggested to play a role in the development of central nervous system, especially the brain. The same CARP X protein are encoded by multiple transcript variants of this gene.

For Research Use Only