

# Recombinant Mouse Carbonic Anhydrase VIII/CA8 Protein (His Tag)

**DIA • AN**<sup>®</sup>  
by Elabscience

Catalog Number: PKSM040495

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	AW546993;Ca8;Cals;Cals1;Car8;Carp;RP23-180H12.1;wdl
<b>Species</b>	Mouse
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met 1-Gln 291
<b>Accession</b>	P28651
<b>Calculated Molecular Weight</b>	34.5 kDa
<b>Observed molecular weight</b>	37 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Measured by its esterase activity. The specific activity is > 5 pmoles/min/μg.

## Properties

<b>Purity</b>	> 88 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 50mM Tris, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 88 % as determined by reducing SDS-PAGE.

## Background

The carbonic anhydrases (or carbonate dehydratases) are classified as metalloenzyme for its zinc ion prosthetic group and form a family of enzymes that catalyze the rapid interconversion of carbon dioxide and water to bicarbonate and protons, a reversible reaction that takes part in maintaining acid-base balance in blood and other tissues. The carbonic anhydrase I (CA) family consists of at least 11 enzymatically active members and a few inactive homologous proteins. Carbonic anhydrase protein (CA) VIII, which is a member of the CA gene family, has been shown to have no catalytic CA activity and its biological function is still unknown. Increased expression of CA-RP VIII was observed in 78% of colorectal carcinomas. It suggested that CA-RP VIII plays a role in the process of invasion in colorectal cancer.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017