## Recombinant Mouse CLEC10A/CD301 Protein (Fc Tag)

### Catalog No. PKSM040532

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	CD301a;M-ASGP-BP-1;Mgl;Mgl1
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Gln 58-Ser 305
Accession	NP_001191181.1
Calculated Molecular Weight	56.7 kDa
Observed molecular weight	58 kDa
Tag	N-hFc
Bioactivity	Not validated for activity
Properties	
Purity	> 86 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	

Data

KDa	МК	R
116	-	
66.2	-	-
45.0	_	
35.0	_	-
25.0	-	
18.4	-	
14.4	-	Sec. 1

> 86 % as determined by reducing SDS-PAGE.

## Background

CLEC10A, also known as the macrophage galactose-type calcium-type lectins (MGLs; CD301) constitute a unique class of C-type lectins because of their specificity for galactose and its structural homologues. MGLs/CD301 is a type II

#### **For Research Use Only**

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

# **Elabscience**®

transmembrane glycoproteins and is expressed on macrophages and related cells of myeloid origins, particularly immature dendritic cells (DCs). There are 2 homologues: MGL1 and MGL2 (CD301a and CD301b) in mice. MGL1/CD301a induces both the production and secretion of interleukin (IL)-10. MGL1/CD301a plays a protective role against colitis by effectively inducing IL-10 production by colonic lamina propria macrophages in response to invading commensal bacteria.

For Research Use Only