

Recombinant Mouse SerpinB12/SERPINB12 Protein (His Tag)

Catalog No. PKSM040612

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	2300003F07Rik;4833409F13Rik
Species	Mouse
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-Pro 423
Accession	NP_082247.1
Calculated Molecular Weight	49.3 kDa
Observed molecular weight	58 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 92 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 100mM NaCl, 0.5mM PMSF, 10% glycerol, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 92 % as determined by reducing SDS-PAGE.

Background

Serpins are the largest and most diverse family of serine protease inhibitors which are involved in a number of

For Research Use Only

fundamental biological processes such as blood coagulation, complement activation, fibrinolysis, angiogenesis, inflammation and tumor suppression and are expressed in a cell-specific manner. Most serpins are secreted and attain physiologic concentrations in the blood and extracellular fluids. Mouse SerpinB12 is a cytoplasm protein which belongs to the serpin family and Ov-serpin subfamily. It is expressed in many tissues, including brain, bone marrow, lymph node, heart, lung, liver, pancreas, testis, ovary, and intestine. SerpinB12 inhibits trypsin and plasmin, but not thrombin, coagulation factor Xa, or urokinase-type plasminogen activator.