

Recombinant Mouse REG3D Protein (His Tag)

Catalog No. PKSM040625

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	INGAP;Ingaprp;MGC130575;Reg3d
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Gly 175
Accession	NP_038921.2
Calculated Molecular Weight	18.5 kDa
Observed molecular weight	17 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

Regenerating islet-derived 3 delta (REG3D) is a member of the secreted Reg superfamily and contains one typical C-type lectin domain. Regenerating gene (Reg), first isolated from a regenerating islet cDNA library, encodes a secretory protein

For Research Use Only

with a growth stimulating effect on pancreatic beta cells. Reg and Reg-related genes which were expressed in various organs have been revealed to constitute a multigene family, the Reg family, which consists of four subtypes (types I, II, III, IV) based on the primary structures of the encoded proteins of the genes, which are associated with tissue repair and have been directly implicated in pancreatic beta-cell regeneration. Reg proteins are expressed in various organs and are involved in cancers and neurodegenerative diseases. They display a typical C-type lectin-like domain but possess additional highly conserved amino acids.