Catalog Number: PKSM040626



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	Complement factor D;28 kDa adipocyte protein;Adipsin;C3 convertase activator;Properdin factor D;Cfd;Adn;Df	
Species	Mouse	
Expression Host	HEK293 Cells	
Sequence	Met 1-Ser 258	
Accession	P03953-2	
Calculated Molecular Weight	27.4 kDa	
Tag	C-His	
Properties		
Purity	> 98 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

KDa	MK	R
116	-	
66.2	-	
45.0	-	-
35.0	-	-
25.0	-	
18.4 14.4	=	
1-4.4	-	

> 98 % as determined by reducing SDS-PAGE.

Background

Complement factor D, also known as Adipsin, C3 convertase activator, Properdin factor D and CFD is a secreted protein which belongs to thepeptidase S1 family. CFD / Adipsin contains onepeptidase S1 domain. Complement factor D (CFD / Adipsin) is a component of the alternative complement pathway best known for its role in humoral suppression of infectious agents. Complement factor D (CFD / Adipsin) has a high level of expression in fat, suggesting a role for adipose tissue in immune system biology. This protein is also a serine protease that is secreted by adipocytes into the bloodstream. Complement factor D (CFD / Adipsin) cleaves factor B when the latter is complexed with factor C3b, activating the C3bbb complex, which then becomes the C3 convertase of the alternate pathway. Its function is homologous to that of C1s in the classical pathway. Complement factor D (CFD / Adipsin) is a serine protease that stimulates glucose transport for triglyceride accumulation in fats cells and inhibits lipolysis. Defects in CFD / Adipsin are

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Catalog Number:PKSM040626



the cause of complement factor D deficiency (CFD deficiency) which predisposes to invasive meningococcal disease.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com