

Recombinant Mouse BAFFR/TNFRSF13C/CD268 Protein (Fc Tag)

Catalog No. PKSM040719

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	2010006P15Rik;BAFF-R;Baffr;Bcmd;Bcmd-1;Bcmd1;Lvis22
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Ala 71
Accession	Q9D8D0-1
Calculated Molecular Weight	33.7 kDa
Observed molecular weight	40-45 kDa
Tag	C-hFc
Bioactivity	Immobilized human BAFF at 10 µg/ml (100 µl/well) can bind mouse BAFFR-Fc, The EC50 of mouse BAFFR-Fc is 0.14-0.32 µg/ml.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Tumor necrosis factor receptor superfamily, member 13C (TNFRSF13C) also known as B-cell-activating factor receptor

For Research Use Only

(BAFFR) and CD268 antigen, is a member of the tumor necrosis factor receptor superfamily. A tumor necrosis factor receptor (TNFR), or death receptor, is a trimeric cytokine receptor that binds tumor necrosis factors (TNF). The receptor cooperates with an adaptor protein which is important in determining the outcome of the response. Members of the TNF receptor superfamily (TNFRSF) have crucial roles in both innate and adaptive immunity and in cellular apoptosis process. Apoptosis is a cell suicide mechanism that enables metazoans to control cell number in tissues and to eliminate individual cells that threaten the animal's survival. Certain cells have unique sensors, termed death receptors or tumour necrosis factor (TNFR), on their surface. Tumour necrosis factors (TNFR) detect the presence of extracellular death signals and, in response, they rapidly ignite the cell's intrinsic apoptosis machinery. It has been proposed that abnormally high levels of BAFFR/TNFRSF13C (CD268) may contribute to the pathogenesis of autoimmune diseases by enhancing the survival of autoreactive B cells.