

Recombinant Mouse SerpinA7/TBG Protein (His Tag)

Catalog No. PKSM040745

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	C730040N12Rik;Tbg
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Gln 418
Accession	NP_808588.3
Calculated Molecular Weight	46.2 kDa
Observed molecular weight	55-60 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

Thyroxine-binding globulin, also known as T4-binding globulin, Serpin A7 and TBG, is a secreted protein which belongs to theserpin family. TBG is synthesized primarily in theliveras a 54 kDa protein.TBG is genomically aserpin, although it

For Research Use Only

has no inhibitory function like many other members of this class of proteins. TBG binds thyroid hormone in circulation. It is one of three proteins (along with transthyretin and albumin) responsible for carrying the thyroid hormones thyroxine (T4) and 3,5,3'-triiodothyronine (T3) in the bloodstream. Of these three proteins, TBG has the highest affinity for T4 and T3, but is present in the lowest concentration. Despite its low concentration, TBG carries the majority of T4 in serum. Due to the very low serum concentration of T4 and T3, TBG is rarely more than 25% saturated with its ligand. Unlike transthyretin and albumin, TBG has a single binding site for T4/T3. TBG tests are sometimes used in finding the reason for elevated or diminished levels of thyroid hormone.