

Recombinant Mouse Prolyl endopeptidase/PREP Protein (His Tag)

Catalog No. PKSM040755

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AI047692;AI450383;D10Wsu136e;PEP;Pop
Species	Mouse
Expression Host	Baculovirus-Insect Cells
Sequence	Leu2-Gln710
Accession	Q9QUR6
Calculated Molecular Weight	82.9 kDa
Observed molecular weight	79-83 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol, 3mM DTT Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Prolyl endopeptidase, also known as PREP, belongs to a distinct class of serine peptidases. It is a large cytosolic enzyme

For Research Use Only

which was first described in the cytosol of rabbit brain as an oligopeptidase. Prolyl endopeptidase degrades the nonapeptide bradykinin at the Pro-Phe bond. It is involved in the maturation and degradation of peptide hormones and neuropeptides such as alpha-melanocyte-stimulating hormone, luteinizing hormone-releasing hormone (LH-RH), thyrotropin-releasing hormone, angiotensin, neurotensin, oxytocin, substance P and vasopressin. Prolyl endopeptidase's activity is confined to action on oligopeptides of less than 10 kD and it has an absolute requirement for the trans-configuration of the peptide bond preceding proline. It cleaves peptide bonds at the C-terminal side of proline residues.