# Recombinant Mouse Prolyl endopeptidase/PREP Protein (His Tag)



Catalog Number: PKSM040755

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

**Synonyms** AI047692;AI450383;D10Wsu136e;PEP;Pop

**Species** Mouse

**Expression Host** Baculovirus-Insect Cells

SequenceLeu2-Gln710AccessionQ9QUR6Calculated Molecular Weight82.9 kDaObserved molecular weight79-83 kDaTagN-His

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per  $\mu g$  of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol, 3mM

DTT


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed m

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

## **Background**

Prolyl endopeptidase, also known as PREP, belongs to a distinct class of serine peptidases. It is a large cytosolic enzyme which was first described in the cytosol of rabbit brain as an oligopeptidase. Prolyl endopeptidase degrades the nonapeptide bradykinin at the Pro-Phe bond. It is involved in the maturation and degradation of peptide hormones and neuropeptides such as alpha-melanocyte-stimulating hormone, luteinizing hormone-releasing hormone (LH-RH), thyrotropin-releasing hormone, angiotensin, neurotensin, oxytocin, substance P and vasopressin. Prolyl endopeptidase's activity is confined to action on oligopeptides of less than 10 kD and it has an absolute requirement for the transconfiguration of the peptide bond preceding proline. It cleaves peptide bonds at the C-terminal side of proline residues.

### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>