

Recombinant Mouse Cathepsin Z Protein (His Tag)

Catalog Number:PKSM040872



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AI787083;AU019819;CTSX;D2Wsu143e
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Val 306
Accession	NP_071720.1
Calculated Molecular Weight	33.2 kDa
Observed molecular weight	38 kDa
Tag	C-His
Bioactivity	Measured by its ability to cleave the fluorogenic peptide substrate, Mca-RPPGFSAFK(Dnp)-OH (R&D Systems, Catalog # ES005). The specific activity is > 1, 200 pmoles/min/μg.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Cathepsin Z (CTS_Z), also known as Cathepsin X or CATX, belongs to the C1 family of lysosomal cysteine proteases. Its gene structure and activity properties show several unique features that distinguish it clearly from other human cysteine proteases. It has a very short pro-region that shows no similarity to those of other cathepsins and a three-residue insertion motif that forms a characteristic 'mini loop'. Cathepsin Z exhibits mono- and di-peptidase activity at its C-terminus, and in contrast to cathepsin B, it does not act as an endopeptidase. It is restricted to the cells of the immune system,

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse Cathepsin Z Protein (His Tag)

Catalog Number:PKSM040872



predominantly monocytes, macrophages and dendritic cells. Cathepsin Z is widely expressed in human tissues, suggesting that this enzyme could be involved in the normal intracellular protein degradation taking place in all cell types. It is capable to cleave regulatory motifs at C-terminus affecting the function of targeted molecules. Cathepsin X may regulate also the maturation of dendritic cells, a process, which is crucial in the initiation of adaptive immunity. Furthermore, higher levels of Cathepsin Z are also found in tumour and immune cells of prostate and gastric carcinomas and in macrophages of gastric mucosa, especially after infection by *Helicobacter pylori*. Cathepsin Z is also ubiquitously distributed in cancer cell lines and in primary tumors from different sources, suggesting that this enzyme may participate in tumor progression.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com