

Recombinant Mouse CD64/FCGR1 Protein (His&AVI Tag)(Active)



Catalog Number:PKSM040878

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	High affinity immunoglobulin gamma Fc receptor I; IgG Fc receptor I; Fc-gamma RI; FcRI; CD64;FcgammaRI;IGGHAFc
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Pro 297
Accession	NP_034316.1
Calculated Molecular Weight	34.3 kDa
Observed molecular weight	45-50 kDa
Tag	C-His-AVI
Bioactivity	Measured by its binding ability in a functional ELISA. Immobilized mouse CD64-AVI at 10 µg/ml (100 µl/well) can bind biotinylated human IgG1, The EC50 of biotinylated human IgG1 is 0.07-0.15 µg/ml.

Properties

Purity	> 97 % as determined by SDS-PAGE
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
Reconstitution	Please refer to the printed manual for detailed information.

Data



Background

High affinity immunoglobulin gamma Fc receptor I, also known as FCGR1 and CD64, is an integral membrane glycoprotein and a member of the immunoglobulin superfamily. CD64 is a high affinity receptor for the Fc region of IgG gamma and functions in both innate and adaptive immune responses. Receptors that recognize the Fc portion of IgG function in the regulation of immune response and are divided into three classes designated CD64, CD32, and CD16. CD64 is structurally composed of a signal peptide that allows its transport to the surface of a cell, three extracellular immunoglobulin domains of the C2-type that it uses to bind antibody, a hydrophobic transmembrane domain, and a short cytoplasmic tail. CD64 is constitutively found on only macrophages and monocytes, but treatment of

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse CD64/FCGR1 Protein (His&AVI Tag)(Active)



Catalog Number:PKSM040878

polymorphonuclear leukocytes with cytokines like IFN γ and G-CSF can induce CD64 expression on these cells. The inactivation of the mouse CD64 resulted in a wide range of defects in antibody Fc-dependent functions. Mouse CD64 is an early participant in Fc-dependent cell activation and in the development of immune responses.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com