

Recombinant Mouse CD5L Protein (His Tag)

Catalog No. PKSM040908

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD5 antigen-like;Apoptosis inhibitor expressed by macrophages;Apoptosis inhibitory 6;CT-2;SP-alpha;Cd5l;Aim;Api6;AAC-11;AI047839;Api6;CT2;Pdp;Sp-alpha
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Val 352
Accession	NP_033820.2
Calculated Molecular Weight	38 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

CD5L, also known as CD5 antigen-like, is a soluble protein belonging to group B of the scavenger receptor cysteine-rich

For Research Use Only

(SRCR) superfamily and contains three SRCR domains. It is a secreted glycoprotein and expressed by macrophages present in lymphoid tissues (spleen, lymph node, thymus, and bone marrow). It binds to myelomonocytic and lymphoid cells and may play an important role in the regulation of the innate and adaptive immune systems. CD5L functions as a pattern recognition molecule by binding both lipoteichoic acid (LTA) on Gram positive and lipopolysaccharide (LPS) on Gram negative bacteria. and the SRCR domain 1 of CD5L retains both the LPS and LTA binding activities. In addition, it is revealed that CD5L seems to play a role as an inhibitor of apoptosis.