A Reliable Research Partner in Life Science and Medicine

# Recombinant Mouse CD27/TNFRSF7 Protein (Fc & His Tag)

Catalog No. PKSM041253

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Synonyms CD27 antigen; CD27L receptor; T-cell activation antigen CD27; Tumor necrosis

factor receptor superfamily member 7;S152;Tnfrsf7;Tp55

Species Mouse

**Expression Host** HEK293 Cells **Sequence** Pro24-Arg182

AccessionP41272Calculated Molecular Weight45.8 kDaObserved molecular weight50-70 kDaTagC-Fc-His

**Bioactivity** Not validated for activity

#### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per μg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



>95~% as determined by reducing SDS-PAGE.

### **Background**

Mouse CD27 is a single-pass type I membrane protein which is a member of the tumor necrosis factor receptor

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

### **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

superfamily. CD27 is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. CD27 transduces signals that lead to the activation of NF-κB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com