

Recombinant Mouse Butyrophilin 1A1/BTN1A1 Protein (His Tag)

Catalog No. PKSM040968

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Butyrophilin subfamily 1 member A1;BTN;BTN1A1;butyrophilin;Btn1a1
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Ala27-Trp247
Accession	Q62556
Calculated Molecular Weight	25.4 kDa
Observed molecular weight	32-35 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Mouse Butyrophilin subfamily 1 member A1 (BTN1A1) is a type I transmembrane glycoprotein which is a member of the Ig superfamily. The BTN1A1 ECD displays two predicted IgV and IgC domains as do B7 and Skint proteins which

For Research Use Only

interact with other Ig superfamily members. BTN1A1 binds to xanthine oxidoreductase (XOR). This interaction stabilizes the association of XOR with the milk fat globule membrane and appears to be essential in the control of milk fat globule secretion. In vitro, BTN1A1 inhibits the proliferation of CD4 and CD8 T-cells activated by anti-CD3 antibodies, T-cell metabolism and IL-2 and IFN- γ secretion. Furthermore, in vivo, BTN1A1 has a protective effect against the development of experimental autoimmune encephalomyelitis (EAE). Because butyrophilins are structurally related to B7 proteins and are functionally implicated in immune regulation, they may represent an emerging family of costimulatory/inhibitory molecules.