

Recombinant Mouse Cadherin-3/CDH3 Protein (Fc Tag)

Catalog No. PKSM040969

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|---|
| Synonyms | Cadherin-3;Placental cadherin;P-cadherin;Cdh3 |
| Species | Mouse |
| Expression Host | HEK293 Cells |
| Sequence | Glu100-Gly647 |
| Accession | P10287 |
| Calculated Molecular Weight | 87.5 kDa |
| Observed molecular weight | 120 kDa |
| Tag | C-Fc |
| Bioactivity | Not validated for activity |

Properties

| | |
|-----------------------|---|
| Purity | > 90 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Cadherin-3 (CDH3) is a single-pass type I membrane protein that belongs to the cadherin superfamily. CDH3 is a calcium-dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region, and a

For Research Use Only

highly conserved cytoplasmic tail. CDH3 is expressed in some normal epithelial tissues and in some carcinoma cell lines. CDH3 preferentially interacts with themselves in a homophilic manner in connecting cells. CDH3 is involved in loss of heterozygosity events in breast and prostate cancer. Mutations in CDH3 have been associated with congenital hypotrichosis with juvenile macular dystrophy.