

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	C-X-C motif chemokine 9;Cxcl9;Gamma-interferon-induced monokine;Monokine induced by interferon-gamma;MIG;Small-inducible cytokine B9;Scyb9
<b>Species</b>	Mouse
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Thr22-Thr126
<b>Accession</b>	AAA39706.1
<b>Calculated Molecular Weight</b>	13.0 kDa
<b>Observed molecular weight</b>	11-17 kDa
<b>Tag</b>	N-His
<b>Bioactivity</b>	Measure by its ability to chemoattract BaF3 cells transfected with mouse CXCR3.The ED <sub>50</sub> for this effect is < 0.3 µg/mL.

## Properties

<b>Purity</b>	> 98 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 0.1 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 98 % as determined by reducing SDS-PAGE.

## Background

Chemokine (C-X-C motif) ligand 9 (CXCL9, MIG), is a small cytokine belonging to the CXC chemokine family. CXCL9 functions as one of the three ligands of chemokine receptor CXCR3 which is a G protein-coupled receptor found predominantly on T cells. It together with CXCL10 and CXCL11, may activate CXCR3 by binding to it. CXCL9 serves as a cytokine that affects the growth, movement, or activation state of cells that participate in immune and inflammatory response. It has been observed that tumour endothelial cells secrete high levels of CXCL9 in all, and CXCL10 in most

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Mouse CXCL9 Protein

Catalog Number:PKSM040997



melanoma metastases. it plays an important role in CD4+ T lymphocyte recruitment and development of CAV, MOMA-2+ macrophages are the predominant recipient-derived source of CXCL9, and recipient CD4 lymphocytes are necessary for sustained CXCL9 production and CAV development in this model.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)