

Recombinant Mouse IGFBP6/IGFBP-6 Protein (His Tag)

Catalog No. PKSM041045

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Insulin-like growth factor-binding protein 6;IBP-6;IGF-binding protein 6;IGFBP-6;Igfbp6;IBP6;IGF binding protein 6;insulin-like growth factor-binding protein 6
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Ala26-Gly238
Accession	P47880
Calculated Molecular Weight	23.7 kDa
Observed molecular weight	28 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Insulin-like growth factors (IGFs) comprise a family of endocrine, paracrine and autocrine polypeptides consisting of the ligands IGF1 and IGF2, two receptors (IGF1R, IGF2R), at least 6 IGF-binding proteins (IGFBPs) and IGFBP proteases. Among the binding proteins, IGFBP6 is unique because of its N-terminal disulfide linkages and its marked binding preference for IGF2. It is a potent inhibitor of the interaction between IGF2 and its receptor IGF1R, thus preventing major functions of IGF2, such as induction of proliferation, differentiation, cell adhesion, or colony formation. In particular, IGFBP-6 inhibited the growth of neuroblastoma and rhabdomyosarcoma xenografts. IGFBP-6 is expressed in many tissues, including lung, liver, gut and the central nervous system.